



## HEPZATO KIT™

(melphalan) for Injection/  
Hepatic Delivery System (HDS)

## HEPZATO KIT: what to ask your care team

It's important to have open communication with your oncologist and healthcare team. This guide suggests questions you may want to ask about treatment with HEPZATO KIT. This guide offers some suggested topics, including

- HEPZATO KIT treatment and procedure
- Potential risks and side effects

This is for informational purposes only and should not be considered medical advice. **Please consult with your physician to see if HEPZATO KIT is right for you.**

### What is HEPZATO and HEPZATO KIT?

**HEPZATO** (melphalan) is a prescription drug for liver-directed treatment for adults with uveal melanoma (a certain type of eye cancer) that has spread to the liver but can't be removed with surgery. This treatment is for cases where less than half of the liver is affected, and the cancer hasn't spread to other parts of the body, or it has only spread to the bones, lymph nodes, skin, or lungs where it can be treated with surgery or radiation.

HEPZATO KIT uses a special delivery system to send the medicine HEPZATO (melphalan) directly to the liver artery and a filtration system to reduce the amount of melphalan in the blood before it goes back into the rest of the body.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about HEPZATO?

- HEPZATO may cause serious problems like bleeding, liver damage, and blood clots. You may need to stop taking blood thinners or blood pressure medicine before treatment with HEPZATO KIT.
- HEPZATO can only be used through a special program called the HEPZATO KIT REMS (Risk Evaluation and Mitigation Strategies) Program.
- HEPZATO can lower the number of blood cells, leading to serious infections, bleeding, or anemia. Tell your doctor right away if you have a fever, bruising, or bleeding.

Please see additional IMPORTANT SAFETY INFORMATION on page 6 and [see here](#) for full Prescribing Information, including Boxed Warning.



What are the risks associated with HEPZATO KIT?

---

---

---

---

---



How is HEPZATO KIT different from other treatments I may have had for my uveal melanoma?

---

---

---

---

---



How long would I have to wait after receiving a different treatment option before I might be eligible for treatment with HEPZATO KIT?

---

---

---

---

---



What additional workups/imagings might be done before my procedure?


---

---

---

---

---



What else should I know about what will happen to me during and after the HEPZATO KIT procedure?

---

---

---

---

---



Will I lose my hair?

---

---

---

---

---



Can filtered blood be put back into my body?

---

---

---

---

---



Are there any dietary or medication restrictions before the procedure?

---

---

---

---

---



How long does it take to recover from the procedure? What type of follow-up is required?

---

---

---

---

---



When can I go back to work? To normal activities?

---

---

---

---

---



Will I still need to take other medicines after I am treated with HEPZATO KIT?

---

---

---

---

---



Why might I need to receive HEPZATO KIT more than once?

---

---

---

---

---



---

---

---

---

---



---

---

---

---

---

[illegible]

# Important Safety Information

## What is HEPZATO and HEPZATO KIT?

HEPZATO (melphalan) is a prescription drug for liver-directed treatment for adults with uveal melanoma (a certain type of eye cancer) that has spread to the liver but can't be removed with surgery. This treatment is for cases where less than half of the liver is affected, and the cancer hasn't spread to other parts of the body, or it has only spread to the bones, lymph nodes, skin, or lungs where it can be treated with surgery or radiation.

HEPZATO KIT uses a special delivery system to send the medicine HEPZATO (melphalan) directly to the liver artery and a filtration system to reduce the amount of melphalan in the blood before it goes back into the rest of the body.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about HEPZATO?

- HEPZATO may cause serious problems like bleeding, liver damage, and blood clots. You may need to stop taking blood thinners or blood pressure medicine before treatment with HEPZATO KIT.
- HEPZATO can only be used through a special program called the HEPZATO KIT REMS (Risk Evaluation and Mitigation Strategies) Program.
- HEPZATO can lower the number of blood cells, leading to serious infections, bleeding, or anemia. Tell your doctor right away if you have a fever, bruising, or bleeding.

### Do not take HEPZATO if you have:

- active brain cancer or brain spots that are likely to bleed
- liver failure, high blood pressure in the liver, or known risk of bleeding in the liver
- surgery or medical treatment of the liver in the past 4 weeks
- blood clotting problems that can't be fixed
- inability to safely undergo general anesthesia, including heart problems like unstable or severe chest pain, heart attack, worsening or new heart failure, serious irregular heartbeats, or severe heart valve disease
- history of allergies or known reactions to melphalan or any part of the HEPZATO KIT, including natural rubber latex, heparin, or severe reactions to iodine contrast not controlled by antihistamines and steroids

### HEPZATO may cause serious side effects including:

**Secondary malignancies**, HEPZATO could have a long-term risk of causing another type of cancer.

**HEPZATO can cause serious birth defects if taken during pregnancy.** Females who can become pregnant should use effective birth control during treatment with HEPZATO and for 6 months after the last dose. Contact your doctor if you become pregnant or think you might be pregnant while

taking HEPZATO. Males with partners who can become pregnant should use effective birth control during treatment with HEPZATO and for 3 months after the last dose.

**Infertility**, HEPZATO has been reported to cause infertility in females and males.

Do not breastfeed during treatment with HEPZATO and for one week after the last dose.

**The most common side effects of HEPZATO are:** low platelets, tiredness, low red blood cells, nausea, muscle and bone pain, low white blood cells, belly pain, vomiting, high liver enzymes, longer blood clotting time, and shortness of breath.

**Please see additional IMPORTANT SAFETY INFORMATION on page 6 and [see here](#) for full Prescribing Information, including Boxed Warning.**

**Call your healthcare provider for medical advice about side effects. To report SUSPECTED ADVERSE REACTIONS, contact Delcath at 1-833-632-0458 and [www.Delcath.com](http://www.Delcath.com) or FDA at 1-800-FDA-1088 or <http://www.fda.gov/medwatch>.**



Copyright © 2025 Delcath Systems, Inc.

All rights reserved.

AMER-P-1506-v1 02/2025

**HEPZATO KIT™**  
(melphalan) for Injection/  
Hepatic Delivery System (HDS)